

An Optical and Near Infrared Search for Brown Dwarfs in the Pleiades Cluster

Chie Nagashima,^{1*} Paul D. Dobbie,² Takahiro Nagayama,¹ Yasushi Nakajima,¹ Tetsuya Nagata,¹ Motohide Tamura,³ Tadashi Nakajima,³ Koji Sugitani,⁴ Hidehiko Nakaya,⁵ Simon T. Hodgkin,⁶ Andrew J. Pickles,⁷ and Shuji Sato¹

¹Department of Astrophysics, Nagoya University, Chikusa-ku, Nagoya 464-8602, Japan

²XROA Group, Department of Physics and Astronomy, University of Leicester, University Road, Leicester LE2 7RH, UK

³National Astronomical Observatory of Japan, Mitaka, Tokyo 181-8588, Japan

⁴Institute of Natural Science, Nagoya City University, Mizuho-ku, Nagoya 467-8501, Japan

⁵Subaru Telescope, National Astronomical Observatory of Japan, Hilo, HI 96720, USA

⁶Institute of Astronomy, University of Cambridge, Madingley Road, Cambridge CB3 0HA, UK

⁷Institute for Astronomy, University of Hawaii, Hilo, HI 96720, USA

Updated on 2003.3.10

ABSTRACT

We have carried out a brown dwarf search over an area of $14' \times 23'$ near the central portion of the Pleiades open cluster in five optical and near infrared bands (i' , Z , J , H , Ks) with 10σ detection limits of $i' \sim 22.0$, $J \sim 20.0$ and $Ks \sim 18.5$ mag. The surveyed area has large extinction in excess of $A_V = 3$ in the Pleiades region. We detected four new brown dwarf candidates from the colour-colour ($J - K$, $i' - J$) and the colour-magnitude (J , $i' - K$) diagrams. We estimated their masses as $0.046 M_\odot$ down to $0.028 M_\odot$. The least massive one is estimated to have a mass smaller than Roque 25 (Martín et al. 1998) or int-pl-IZ-69 (Dobbie et al. 2002b), and maybe the lowest mass object found so far in the Pleiades cluster.

Key words: open clusters and associations: individual (Pleiades) – stars: low-mass, brown dwarfs

1 INTRODUCTION

Many brown dwarf (BD) searches have targeted young open clusters or star forming regions rather than general fields, because here BDs are still relatively bright and warm during their early phase. The Pleiades region is one of the best clusters for this purpose; it is a fairly young, nearby, rich and compact cluster. It has an age of 125 Myr (Stauffer, Schultz, & Kirkpatrick 1998) at a distance of 130 pc (Crawford & Perry 1976; Pinfield et al. 2000). Twelve hundred member stars are located within 2.5 degrees of the cluster centre (Pinfield, Jameson, & Hodgkin 1998).

A number of optical surveys searching for BDs in the Pleiades cluster have been done to date and identified a numerous population of bona-fide BDs and BD candidates; “CFHT-PL” objects found by an RI survey (Bouvier et al. 1998); “NPL” objects by an $RIJK$ survey (Festin 1998); “MHObd” objects by a VI survey

(Stauffer et al. 1998b); “IPMBD” objects by an RI survey (Hambly et al. 1999); “Roque” objects by an IZ survey (Zapatero Osorio et al. 1999); “BPL” objects by an IZ survey (Pinfield et al. 2000); “int-pl-IZ” objects by an IZ survey (Dobbie et al. 2002b). About 20 of these objects are confirmed as Pleiades brown dwarfs on the basis of observed lithium abundance, radial velocity and proper motion (e.g., Rebolo, Zapatero-Osorio, & Martín 1995; Rebolo et al. 1996; Basri, Marcy, & Graham 1996; Zapatero Osorio et al. 1997a; Stauffer, Schultz, & Kirkpatrick 1998; Martín et al. 2000). However, to date, only one L-type BD candidate (Roque25; Martín et al. 1998) has been spectroscopically identified in the Pleiades cluster. Detection of L-type BDs is difficult at optical wavelengths because they should emit most of their bolometric flux in the infrared region. We have carried out a brown dwarf survey in both the optical and near infrared of a small region of the central region of the Pleiades cluster or southern portion of Merope. Most existing surveys have ignored this region because of its proximity to a small molecular cloud which results in large and variable extinction (Breger 1987).

* E-mail:chie@optik.mtk.nao.ac.jp; Present address: National Astronomical Observatory of Japan, Mitaka, Tokyo 181-8588, Japan




Figure 1. The field locations on the sky. The shaded square shows the observed area and the open squares show the region covered only in optical observation. The overplotted dashed circles are radii of 0.26° and 0.64° from the centre of the cluster, R.A. 3^h47^m , Dec. $+24^\circ07'$ (J2000.0; Pinfield et al. 2000).

2 OBSERVATIONS

We surveyed an area of $14' \times 23'$ centred at R.A. $3^h46^m59^s.6$, Dec. $+23^\circ44'59''$ (J2000.0) both in the optical and near infrared bands. The observed area is shown as the shaded rectangle in Fig. 1. The four open rectangles show the area covered at optical wavelengths.

2.1 Near Infrared Survey

Our near infrared survey was made on 2000 October 12, 17 and 18 with SIRIUS (Simultaneous InfraRed Imager for Unbiased Survey) on the University of Hawaii 2.2 m telescope, atop Mauna Kea. SIRIUS employs three 1024×1024 pixel HgCdTe arrays which provides J ($1.25\mu\text{m}$), H ($1.65\mu\text{m}$), and Ks ($2.15\mu\text{m}$) band images simultaneously (Nagashima et al. 1999; Nagayama et al. 2002). The field of view in the each band is 4.9×4.9 with a pixel scale of $0''.28$ at the Cassegrain focus of $f/10$. We mapped an area of $14' \times 23'$ with 3×5 tiles of the field of view. We obtained each of the tiles by dithering 18 frames. The exposure time was 60 sec for each frame. The limiting magnitudes at 10σ are 20.0, 19.2, and 18.5 at J , H , and Ks , respectively. Typical seeing in the Ks band was $1''.0$.

Dark and dome-flat frames were taken at the beginning or the end of each night. We observed standard stars, Nos. 9105, 9107, 9108, 9116 and 9188, in the faint NIR standard star catalog of Persson et al. (1998) on the same nights for photometric calibration, and the red standard stars, LDN 547, BRI B0021-0214 and BRI B2202-1119, in Table 3 of Persson et al. (1998) on 2001 August 31 for determination of the colour conversion formula from ours to the CIT system. We obtained the following equations as colour-conversion from ours to the CIT system.

$$J_{\text{CIT}} = J_{\text{obs}} - (0.013 \pm 0.017) \times (J - Ks)_{\text{obs}} + (0.012 \pm 0.030)$$

$$H_{\text{CIT}} = H_{\text{obs}} - (0.029 \pm 0.029) \times (J - H)_{\text{obs}} + (0.007 \pm 0.033)$$

$$K_{\text{CIT}} = Ks_{\text{obs}} - (0.024 \pm 0.020) \times (J - Ks)_{\text{obs}} + (0.018 \pm 0.035)$$

We used pipeline software based on NOAO's IRAF

(Imaging Reduction & Analysis Facility)¹ package to reduce the data. We have applied standard NIR image reduction procedures, including dark current subtraction, sky subtraction and flat fielding. Source detection, morphological classification and photometry were performed with the APPHOT package in IRAF.

2.2 Optical Survey

The i' ($0.77\mu\text{m}$, SDSS system) and Z ($0.9\mu\text{m}$, RGO) band image was taken on 2000 October 20 with the Wide Field Camera (WFC) on the Isaac Newton Telescope (INT) at La Palma, as part of the INT Wide Angle Survey (McMahon et al. 2001). The WFC accommodates four 2048×4196 pixel EEV CCDs. The field of view for each CCD is 11.4×22.8 with a pixel scale of $0''.33$ at the prime focus of the INT and the mosaic layout of the four CCDs enables us to encompass the whole field of $14' \times 23'$ taken in the near infrared bands (Fig. 1). We obtained one frame for each filter; the exposure time of each frame was 600 s. The 10σ detection limits are 22.0 and 20.5 at i' and Z , respectively, where i' and Z are on natural system of WFC.

The observed data were reduced at the Institute of Astronomy, Cambridge, using the WFC data reduction pipeline software (Irwin & Lewis 2001). All the frames were bias-subtracted and then corrected for linearity before the science fields were flatfielded and deFRINGED. The pipeline was also used to generate photometrically and astrometrically calibrated object catalogues. On the night in question, the i' - and Z -band photometric zeropoints were measured from three standard star fields (observed once each) in the catalogue of Landolt (1992) and were found to be stable to about 1 %. The astrometric solution takes account of the radial distortion across the WFC field of view, and resulted in residuals of $0''.22$.

3 SELECTION OF PLEIADES BD CANDIDATES

3.1 The colour-colour diagram

We constructed the colour-colour ($J - K$, $i' - J$) diagram for all the objects, 846 sources in total, detected with S/N ratios higher than 10 in the near infrared and optical bands, differentiating them as extended (149; open circles) and point (697; filled circles) sources in Fig. 2. Seven sources detected with S/N ratios higher than 10 in the near infrared but lower than 5 in the optical are also plotted with large error bars. "Extended objects" were sorted with a criterion of the FWHMs larger by at least 20 % than the seeing size of the night in the Ks images.

The thick solid and dashed curves from the bottom-left corner represents the loci of main sequence and red giant stars (Bessell 1991, 1979); spectral types of K7 dwarf at the cusp at around (1,0.8) and of M7 at the tip (3.5,1.2) for ($i' - J$, $J - K$), while the arrow at the upper-right corner shows

¹ IRAF is distributed by the National Optical Astronomy Observatory, which are operated by the Association of Universities for Research in Astronomy, Inc., under cooperative agreement with the National Science Foundation.




Figure 2. The $J - Ks, i' - J$ colour-colour diagram for all objects detected in all the 5 bands. Small dots are point sources and open circles are extended sources. Asterisks are the known late-type field dwarfs. The thick solid and dashed curves from the bottom left corner represent the loci of main sequence and red giant stars. The arrow at the upper right corner shows the reddening vector of $A_V = 3$ mag. The thin dashed line is the ~ 125 Myr isochrone of the NEXTGEN model (Baraffe et al. 1998), and the thin dot-dashed line is the ~ 120 Myr isochrone of the DUSTY model (Chabrier et al. 2000). Numbers with tick marks denote masses in solar mass units for each model.

the reddening vector of $A_V = 3$ mag (Schultz & Wiemer 1975; Rieke & Lebofsky 1985). Most of the point sources (filled circles) are concentrated above the locus from K7 to M7, along the extinction vector up to 3 mag from the main sequence (reddening band), indicating they are K- or M-type dwarfs lying intrinsically on the main sequence. On the other hand, most of the extended sources and some point sources spread upward of the reddening band. About 20 of the extended sources have galaxy-like shapes. The K-correction curves from $z = 0 - 0.1$ of elliptical and spiral galaxies (Furusawa, private communication) are plotted as dotted lines. Thus, objects spread upward of the reddening band includ-

ing the point sources (182 objects in total) are likely to be elliptical or spiral type galaxies reddened with K-corrections of $z \sim 0$ to 0.1 and A_V of 0 to several magnitudes. A major problem in this kind of survey is to distinguish stars from distant galaxies at faint magnitudes (e.g. Festin 1997). The optical and near-infrared colour-colour diagram allows us to easily distinguish them. Note that the near infrared colour-colour diagram ($J - H, H - K$) alone cannot distinguish low-mass stars from these galaxies, due to merging of them in the diagram (e.g. Kirkpatrick et al. 1999; Jarrett et al. 2000).

To highlight the expected location of Pleiades BDs in

the colour-colour diagram, two theoretical isochrones for the Pleiades cluster are also plotted in Fig. 2; the dashed line is the 125 Myr isochrone of the NEXTGEN model (Baraffe et al. 1998), while the dot-dashed line is the 120 Myr isochrone of the DUSTY model (Chabrier et al. 2000, ; Baraffe, private communication for 120 Myr model). The I_C magnitudes of these models have been transformed on to our system using the following relation, $i' = I_C + 0.211(R_C - I_C) + 0.011(V - I_C)^2 - 0.003(V - I_C)$. This transformation is derived via Landolt system (I_{Lan}); the transformation between i' and I_{Lan} is from an INT web page ², and that between I_{Lan} and I_C is from Bessell (1990). The NEXTGEN model is based on the nongray dust-free atmosphere (Hauschildt, Allard, & Baron 1999), and describes successfully various observed properties of early-mid M dwarfs, but predicts near infrared colours too blue compared to the observed ones for cooler objects. The DUSTY model includes the effect of grain formation, that is, 1) the photospheric depletion of dust-forming elements, and 2) scattering and absorption by dust, explaining successfully the near infrared colours of the late M and L types with $1,800 \lesssim T_{\text{eff}} \lesssim 2,200$ K (e.g. Jameson et al. 2002). Because of the inadequacy of the NEXTGEN model in the cooler BD region, we have adopted the DUSTY model in the following discussion. Using near infrared photometry and lithium tests of low mass cluster members, Zapatero Osorio et al. (1997b) estimated the stellar / substellar boundary to be located at an absolute magnitude of ($\sim 0.075 M_{\odot}$) for Pleiades (~ 120 Myr) to be at $M_I = 12.4$, $M_J = 10.0$ and $M_K = 9.0$ mag. This corresponds to the point $(i' - J, J - K) = (2.9, 1.0)$ in our colour-colour diagram, highlighted by a cross, at approximately spectral type M6. Twelve late-type field dwarfs from Dobbie et al. (2002a) were also plotted as asterisks. We adopt the redder side of the line running from the cross in parallel with the extinction vector as the location of Pleiades BDs in Fig. 2. Note that this location coincides with that of late-M and L-type field dwarfs. There remain 12 sources on the redder side of the line.

3.2 The colour-magnitude diagram

To separate Pleiades BDs from field dwarfs, we construct the colour-magnitude diagram ($J, i' - K$). The 12 red objects selected by Fig. 2 are plotted in Fig. 3. The theoretical tracks representative of the low mass stellar and substellar members of the Pleiades are also plotted; NEXTGEN model (Baraffe et al. 1998) and DUSTY model (Chabrier et al. 2000). We choose the J magnitude as the ordinate of the diagram because i' magnitudes of the red objects have larger errors and K magnitudes are less sensitive to the mass difference on the theoretical tracks. The arrow at the upper-right corner indicates the reddening vector of $A_V = 3$ mag, which runs nearly in parallel with the tracks.

We adopt the distance modulus used by Dobbie et al. (2002b), $(m - M)_0 = 5.53$, to estimate the ordinate displacement (J magnitude) of the isochrone. Different age estimates (70 – 150 Myr) result in a displacement of ~ -0.3 and $\sim +0.1$ mag. Different distance estimates result in a displacement of ~ -0.2 mag (*Hipparcos*) and $\sim +0.1$ mag (photometry). The

effect of the cluster depth results in a displacement of $\sim \pm 0.2$ mag. Additionally, location of unresolved binaries is by 0.75 mag above the single-star sequence (e.g. Pinfield et al. 2000). Accounting for all of these uncertainties and allowing for a small degree of error in both of the theoretical models and photometry, BDs belonging to the Pleiades cluster should be located between 0.3 mag below and 1.0 mag above the relevant single-star isochrone.

With these criteria, we identified four Pleiades BD candidates from 12 red objects. The remaining eight objects lie more than 1 mag below the isochrone in the colour-magnitude diagram, although they fall in the Pleiades BD domain in the colour-colour diagram. They could be late-M and L-type dwarfs lying background of Pleiades. Finally, we list the four sources (PL- # of Fig. 3) as BD candidates from our five-colour survey. All of them are previously unpublished. The positions and magnitudes of four BD candidates are provided in Table 1, and that of the remaining eight objects are in Table 2. Finding charts of four BD candidates are presented in Fig. 4.

4 DISCUSSION

4.1 Estimation of the BD candidate masses

To estimate masses of BD candidates from their observed magnitudes, we should determine the level of interstellar extinction towards them. The distribution of the background field stars in Fig. 2 indicates that the interstellar extinction in the surveyed portion of Pleiades ranges from 0 to 3 mag in A_V . The foreground extinction toward Pleiades is relatively uniform with the value of $A_V \sim 0.12$ mag, except for a region south of Merope, which contains a small molecular cloud (Crawford & Perry 1976). Because the cloud is assigned to be inside the cluster from polarimetry and reddening of the member stars and the field stars (Bregier 1987), the extinctions estimated from background stars should be regarded as upper limit for the BD candidates.

To estimate the maximum interstellar extinction for the individual BD candidates, we constructed an extinction map in the surveyed area from all the field stars (small dots between the two straight lines in Fig. 2), assuming them to be K- or M-type main sequence stars (the thick line in Fig. 2). Visual extinction, A_V , toward each star was calculated from its $i' - J$ and $J - K$ colours with the reddening law of Schultz & Wiemer (1975) and Rieke & Lebofsky (1985), i.e. $E(J - K)/E(i' - J) = 0.56$, $A_V = 6.10 \times [(J - K)_{\text{obs}} - (J - K)_{\text{intrinsic}}]$. We meshed the area with 17×10 grids of $1'$ square and adopted the maximum A_V within the grid, and averaged them over each mesh with Gaussian weights. The final A_V map is shown in Fig. 5 and the values toward the BD candidates are listed in Table 3.

The estimated maximum extinctions for individual candidates are shown in Fig. 3 with dotted arrows. If extinction is neglected, PL-2 and -3 are estimated to have $M \sim 0.035 M_{\odot}$, similar to Roque 25 (Martín et al. 1998) and int-pl-IZ-69, -81, -84 (Dobbie et al. 2002b), the coolest candidate members so far detected. On the other hand, assuming the maximum likely level of extinction, their masses are estimated to be $M \sim 0.04 M_{\odot}$. Similarly we estimate the mass of the faintest candidate to lie in the range of $0.033 - 0.028$

² <http://www.ast.cam.ac.uk/~wfcsur/photom.php>




Figure 3. The $J, i' - K$ colour-magnitude diagram. The BD candidates are shown with filled circles. Dotted arrow from each candidates shows the estimated maximum reddening vector (see Section 4.1). Other symbols are as in Fig. 2. The $i' - K$ colour of Roque 25 (Martín et al. 1998) is also shown with a perpendicular line due to lack of J magnitude.

Table 1. The positions and magnitudes of new brown dwarf candidates.

ID	R.A.	Dec.	i'	Z	J	H	K	name
c06	03:45:50.6	+23:44:37	20.02 ± 0.02	18.19 ± 0.01	16.34 ± 0.06	15.68 ± 0.07	15.19 ± 0.07	PL-1
c08	03:46:34.3	+23:50:04	21.36 ± 0.06	19.52 ± 0.03	17.35 ± 0.06	16.63 ± 0.07	16.01 ± 0.08	PL-2
c10	03:45:11.7	+23:41:44	21.98 ± 0.09	19.88 ± 0.05	17.59 ± 0.06	16.79 ± 0.07	16.13 ± 0.08	PL-3
c12	03:45:58.5	+23:41:54	$23.40^{+0.40}_{-0.29}$	$21.69^{+0.29}_{-0.23}$	18.54 ± 0.08	17.42 ± 0.09	16.53 ± 0.09	PL-4

Table 2. The positions and magnitudes of other red objects.

ID	R.A.	Dec.	i'	Z	J	H	K
c01	03:46:24.9	+23:46:10	$22.28^{+0.12}_{-0.11}$	$20.93^{+0.13}_{-0.12}$	19.30 ± 0.08	18.90 ± 0.12	18.28 ± 0.13
c02	03:45:30.9	+23:49:51	22.03 ± 0.1	20.45 ± 0.08	18.97 ± 0.07	18.32 ± 0.08	17.91 ± 0.09
c03	03:45:59.7	+23:40:53	22.33 ± 0.14	20.82 ± 0.12	18.93 ± 0.07	18.35 ± 0.08	17.89 ± 0.10
c04	03:46:26.6	+23:39:35	21.57 ± 0.07	20.14 ± 0.06	18.01 ± 0.06	17.22 ± 0.07	16.72 ± 0.08
c05	03:46:28.0	+23:42:39	22.19 ± 0.12	21.11 ± 0.15	18.60 ± 0.07	17.74 ± 0.08	17.29 ± 0.09
c07	03:46:20.7	+23:41:56	$23.18^{+0.35}_{-0.27}$	$22.18^{+0.52}_{-0.35}$	19.14 ± 0.08	18.75 ± 0.10	18.03 ± 0.11
c09	03:45:50.9	+23:44:57	$22.65^{+0.19}_{-0.16}$	$20.84^{+0.13}_{-0.12}$	18.63 ± 0.07	17.64 ± 0.08	17.08 ± 0.09
c11	03:45:49.2	+23:41:26	$24.34^{+1.49}_{-0.61}$	$21.93^{+0.26}_{-0.21}$	19.88 ± 0.10	18.78 ± 0.13	18.33 ± 0.14




Figure 4. Finding charts for new brown dwarf candidates (K_s band). Each panel is $2' \times 2'$, North is up, East is left.




Figure 5. The extinction map constructed from the $J - K$ and $i' - J$ colours of the point sources in the reddening band. Small dots are field stars and diamonds are BD candidates. Estimation of the extinction is described in Section 4.1. The correspondence between grey scale and A_V is indicated in the scale bar.

$M_{\odot}$. Therefore, PL-4 maybe the lowest mass member of the Pleiades identified to date. The estimated masses from J magnitudes for the case of no extinction and that of maximum possible extinction toward individual candidates are shown in Table 3. Note that uncertainties in the age and the distance of the cluster and the effect of the cluster depth contribute an additional error of 10 – 20 % in the mass estimation; e.g. the mass of PL-4 for the case of no extinction is $0.028^{+0.002}_{-0.001} M_{\odot}$. We note that our mass estimates would increase slightly if we were to take into account the “missing M dwarf gap” (Dobbie et al. 2002c).

Table 3. Mass estimates for brown dwarf candidates based on 120 Myr DUSTY model.

Name	MAX A_V	Mass($M_{\odot}$)	
		$A_V=0 - \text{MAX}$	SpT
PL-1	2.5	0.046 – 0.063	M6 – M8
PL-2	2.0	0.035 – 0.040	M8 – L1
PL-3	2.5	0.033 – 0.039	M8 – L1
PL-4	3.5	0.028 – 0.033	L1 – L5




Figure 6. The broad-band energy distributions for four brown dwarf candidates. Solid and dashed lines represent brown dwarf candidates with no extinction and with possible maximum extinction, respectively. M6 – M8 and L0 – L5 field dwarfs are overplotted with dotted lines.

4.2 Broad-band energy distributions

In Fig. 6 we compare the four SEDs with those of cool field dwarfs, determined in previous surveys (M6 – M8; Kirkpatrick & McCarthy 1994, L0 – L5; Kirkpatrick et al. 1999, 2000). Solid and dashed lines represent SEDs of the four BD candidates for the cases of no extinction and maximum extinction discussed in § 4.1, respectively. We have assumed the distance modulus for Pleiades of $(m - M)_0 = 5.53$ (Crawford & Perry 1976; Pinfield et al. 2000) to convert the observed magnitudes to absolute magnitudes for the cluster member. The zero-magnitude fluxes are taken from Bessell, Castelli, & Plez (1998).

All the candidates show a steep decline towards the shorter wavelengths and a broad hump between 1 and 2 μm . Within the uncertainties arising from the correction for interstellar extinction, the four SEDs of the candidate BDs are well matched by the energy distributions of the late-M and L-type field dwarfs: PL-1 is consistent with a mid to late-M spectral type, PL-2 and PL-3 with late-M to early L spectral type and PL-4, an early-mid L spectral type. We summarize these results in Table 3.”

4.3 Contamination of field stars

As discussed in section 3.1, the optical and near-infrared colour-colour diagram allows us to separate distant galaxies and giants from BD candidates. However, field dwarfs are still possible sources of contamination.

The DUSTY model indicates that Pleiades BDs are over-luminous with respect to field stars of similar spectral type by ~ 1 mag. The criterion of selecting objects between 0.3 mag below and 1.0 mag above the isochrone means that we are sensitive to single and binary dwarfs at distance of 71 – 130 and 96 – 184 pc, respectively. This corresponds to space volumes of 15.5 and 45.1 pc³ for single and binary field stars, respectively. Assuming a space density of M8 – L4.5 dwarfs of 0.0066 pc⁻³ (Gizis et al. 2000) and a binary fraction of 50 % (e.g. Steele & Jameson 1995), we estimate the contamination of ~ 0.2 field star in our BD candidates. Thus, the level of the contamination of the BD candidates by non-members is negligibly low.

4.4 The cluster luminosity function and mass function

We have derived the cluster luminosity function for J magnitudes between 15.5 to 20.0 which correspond to the BD limit of Zapatero Osorio et al. (1997b) and the 90 % completeness limit of our survey (almost the same as 10 σ detection limit), respectively. In our calculation we assume that the interstellar extinction toward Pleiades in our survey area is spatially uniform with the value of 1.5 mag at V or 0.4 mag at J , we subdivided this magnitude range into 2-mag bins, given in Table 4.

Using the DUSTY model, we have calculated the masses corresponding to each of our luminosity bins. We estimated the total number, N_{annul} , of member of the whole cluster from the number of our sample. First we calculate the number, N_{annul} , of members inside an annulus. Our survey area covers 7.7 % of the area of the annulus of an inner and an outer radius of $r_i = 0.26^\circ$ and $r_o = 0.64^\circ$, respectively (Fig 1). N_{annul} is calculated simply from the area ratio. Next we calculate N_{total} . Pinfield, Jameson, & Hodgkin (1998) and Raboud & Mermilliod (1998) show that the surface density distribution of the Pleiades members can be well fitted by a King distribution (King 1962) whose core radius increases as the stellar mass decreases. The following equation, which is obtained by integrating the function of King distribution, provides the number of cluster members inside the circle of a radius r ,

$$n(r) = \pi r_c^2 k \left[\ln(1+x) - 4 \frac{\sqrt{1+x} - 1}{\sqrt{1+x_t}} + \frac{x}{1+x_t} \right], \quad (1)$$

where $x = (r/r_c)^2$, $x_t = (r_t/r_c)^2$, k is the surface density of the cluster, r the radius from the cluster centre, r_c the core radius at which the surface density of members falls to half its central value, and r_t the tidal radius at which the external gravitational potential equals the cluster potential. The total number of cluster members, N_{total} , can be determined independently of k by taking the ratio $n(r_t)/[n(r_o) - n(r_i)]$, where $n(r_o) - n(r_i)$ is equal to N_{annul} and $n(r_t)$ is equal to N_{total} . We adopt a value of $r_c = 2.2^\circ$ from Jameson et al. (2002) for low-mass members of Pleiades, and a value of $r_t = 5.54^\circ$ from Pinfield, Jameson, & Hodgkin (1998).

$$\frac{N_{\text{total}}}{N_{\text{annul}}} = \frac{n(r_t)}{[n(r_o) - n(r_i)]} = 12.78. \quad (2)$$

Finally, we calculate the values of substellar mass function of the cluster, N_{total} per unit mass. The values are given in Table 4 and are plotted in Fig. 7 with several previous estimates (Dobbie et al. 2002b; Hodgkin & Jameson 2000; Hambly et al. 1999). The shape of mass function of the Pleiades across and below the stellar / substellar boundary has been reported to be well represented by a power-law with an index α lying between 0.4 – 1.0, dependent on investigator (e.g. Jameson et al. 2002; Bouvier et al. 1998; Hodgkin & Jameson 2000; Zapatero Osorio et al. 1997a). Our new points seem to support the larger values of α . However, as the present survey covers only a small fraction of the cluster (i.e., $< 0.5\%$), this result must be treated with a measure of caution. Nevertheless, we note that Béjar et al. (2001) have recently determined the substellar mass function of the much younger σ Orionis cluster to be well matched by a power-law with index $\alpha = 0.8$.

5 CONCLUSIONS

We have undertaken an optical (i' , Z) and near infrared (J , H , K_s) survey of a $14' \times 23'$ area near the centre of the Pleiades cluster to search for brown dwarf members. We have unearthed 4 BD candidates (PL-1, -2, -3, and -4), out of 853 sources. PL-1 is likely an M-type BD, PL-2 and -3 are likely late-M or early-L type BDs, while PL-4 is probably an L-type BD. The estimated mass of our faintest candidate, PL-4, is in the range of 0.028 – 0.033 $M_\odot$, lower than that of Roque 25 (Martín et al. 1998) and can be lower than that of int-pl-IZ-69 (Dobbie et al. 2002b), possibly making it lowest mass Pleiad identified to date.

ACKNOWLEDGMENTS


We thank the staffs of the UH 2.2 m telescope and the Subaru telescope for supporting the observation of SIRIUS. This work is financially supported by SUMITOMO foundation, Grant-in-Aid for Scientific Researches on Priority Areas (A), and International Scientific Research, the Ministry of Education, Culture, Sports, Science, and Technology. The Isaac Newton Telescope is operated by the Isaac Newton Group in the Spanish Observatorio del Roque de los Muchachos of the Instituto de Astrofísica de Canarias. Nagashima, Nagayama, and Y. Nakajima are financially supported by the Japan Society for the Promotion of Science. PDD is a PPARC postdoctoral research associate. We also thank an anonymous referee for his/her useful comments and advice.

REFERENCES

- Baraffe, I., Chabrier, G., Allard, F., & Hauschildt, P. H. 1998, A&A, 337, 403
- Baraffe, private communication
- Basri, G., Marcy, G. W., & Graham, J. R. 1996, ApJ, 458, 600
- Béjar, V. J. S. et al. 2001, ApJ, 556, 830
- Bessell, M. S. 1979, PASP, 91, 589

Table 4. N_{total} per unit mass (in the case of $A_V = 1.5$ mag).

J magnitude bin	Mass bin ($M_{\odot}$)	Mid-mass ($M_{\odot}$)	N_{obs}	N_{annul}	N_{total}	N_{total} per unit mass
15.1 – 17.1	0.095 – 0.037	0.066	2	26	330	$5,700 \pm 4,000$
17.1 – 19.6	0.037 – 0.025	0.031	2	26	330	$28,000 \pm 19,000$

**Figure 7.** The mass function in the Pleiades cluster. Open stars represent this work. Filled stars, squares, and circles represent the results of Dobbie et al. (2002b), Hodgkin & Jameson (2000), and Hambly et al. (1999), respectively, together with the model fits by Hodgkin & Jameson (2000) (solid line), Hambly et al. (1999) (dashed line), van Leeuwen (1980) (dot-dashed line).

Bessell, M. S. 1990, PASP, 102, 1181

Bessell, M. S. 1991, AJ, 101, 662

Bessell, M. S., Castelli, F., & Plez, B. 1998, A&A, 333, 231

Bouvier, J., Stauffer, J. R., Martin, E. L., Barrado y Navascués, D., Wallace, B., & Bejar, V. J. S. 1998, A&A, 336, 490

Breger, M. 1987, ApJ, 319, 754

Crawford, D. L. & Perry, C. L. 1976, AJ, 81, 419

Chabrier, G., Baraffe, I., Allard, F., & Hauschildt, P. 2000, ApJ, 542, 464

Dobbie, P. D., Kenyon, F., Jameson, R. F., & Hodgkin, S. T. 2002a, MNRAS, 331, 445

Dobbie, P. D., Kenyon, F., Jameson, R. F., Hodgkin, S. T., Pinfield, D. J., & Osborne, S. L. 2002b, MNRAS, 335, 687

Dobbie, P. D., Pinfield, D. J., Jameson, R. F., & Hodgkin, S. T. 2002c, MNRAS, 335, L79

Festin, L. 1997, A&A, 322, 455

Festin, L. 1998, A&A, 333, 497

Furusawa, private communication

Gizis, J. E., Monet, D. G., Reid, I. N., Kirkpatrick, J. D., Liebert, J., & Williams, R. J. 2000, AJ, 120, 1085

Hambly, N. C., Hodgkin, S. T., Cossburn, M. R., & Jameson, R. F. 1999, MNRAS, 303, 835

Hauschildt, P. H., Allard, F., & Baron, E. 1999, ApJ, 512, 377

Hodgkin, S. T. & Jameson, R. F. 2000, ASP Conf. Ser. 198: Stellar Clusters and Associations: Convection, Rotation, and Dynamos, 59

Irwin, M. & Lewis, J. 2001, New Astronomy Review, 45, 105

Jameson, R. F., Dobbie, P. D., Hodgkin, S. T., & Pinfield, D. J. 2002, MNRAS, 335, 853

Jarrett, T. H., Chester, T., Cutri, R., Schneider, S., Skrutskie, M., & Huchra, J. P. 2000, AJ, 119, 2498

King, I. 1962, AJ, 67, 471

Kirkpatrick, J. D. & McCarthy, D. W. 1994, AJ, 107, 333

Kirkpatrick, J. D. et al. 1999, ApJ, 519, 802

Kirkpatrick, J. D. et al. 2000, AJ, 120, 447

Landolt, A. U. 1992, AJ, 104, 340

Martín, E. L., Basri, G., Zapatero-Osorio, M. R., Rebolo, R., & López, R. J. G. 1998, ApJ, 507, L41

Martín, E. L., Brandner, W., Bouvier, J., Luhman, K. L., Stauffer, J., Basri, G., Zapatero Osorio, M. R., & Barrado y Navascués, D. 2000, ApJ, 543, 299

McMahon, R. G., Walton, N. A., Irwin, M. J., Lewis, J. R., Bunclark, P. S., & Jones, D. H. 2001, New Astronomy Review, 45, 97

Nagashima, C. et al. 1999, Star Formation 1999, Proceedings of Star Formation 1999, held in Nagoya, Japan, June 21 - 25, 1999, Editor: T. Nakamoto, Nobeyama Radio Observatory, p. 397-398, 397

Nagayama, T. et al. 2002, Proceedings of the SPIE, 4841, in press

Persson, S. E., Murphy, D. C., Krzemiński, W., Roth, M., & Rieke, M. J. 1998, AJ, 116, 2475

Pinfield, D. J., Jameson, R. F., & Hodgkin, S. T. 1998,

- MNRAS, 299, 955
- Pinfield, D. J., Hodgkin, S. T., Jameson, R. F., Cossburn, M. R., Hambly, N. C., & Devereux, N. 2000, MNRAS, 313, 347
- Raboud, D. & Mermilliod, J.-C. 1998, A&A, 329, 101
- Rebolo, R., Zapatero-Osorio, M. R., & Martin, E. L. 1995, Nature, 377, 129
- Rebolo, R., Martin, E. L., Basri, G., Marcy, G. W., & Zapatero-Osorio, M. R. 1996, ApJL, 469, L53
- Rieke, G. H. & Lebofsky, M. J. 1985, ApJ, 288, 618
- Schultz, G. V. & Wiemer, W. 1975, A&A, 43, 133
- Stauffer, J. R., Schultz, G., & Kirkpatrick, J. D. 1998, ApJL, 499, L199
- Stauffer, J. R., Schild, R., Barrado y Navascues, D., Backman, D. E., Angelova, A. M., Kirkpatrick, J. D., Hambly, N., & Vanzi, L. 1998, ApJ, 504, 805
- Steele, I. A. & Jameson, R. F. 1995, MNRAS, 272, 630
- van Leeuwen, F. 1980, IAU Symp. 85: Star Formation, 85, 157
- Zapatero Osorio, M. R., Rebolo, R., Martin, E. L., Basri, G., Magazzu, A., Hodgkin, S. T., Jameson, R. F., & Cossburn, M. R. 1997a, ApJL, 491, L81
- Zapatero Osorio, M. R., Martin, E. L., & Rebolo, R. 1997b, A&A, 323, 105
- Zapatero Osorio, M. R., Rebolo, R., Martín, E. L., Hodgkin, S. T., Cossburn, M. R., Magazzù, A., Steele, I. A., & Jameson, R. F. 1999, A&AS, 134, 537

This paper has been typeset from a $\text{\TeX}$ / $\text{\LaTeX}$ file prepared by the author.